

цинской ассоциации о гуманном обращении с животными, 1996 г.).

КОНФЛИКТ ИНТЕРЕСОВ

Авторы заявляют, что у них нет конфликта интересов.

СПИСОК ЛИТЕРАТУРЫ

- Калинина Н.И., Сысоева В.Ю., Рубина К.А., Парфенова Е.В., Ткачук.* 2011. Мезенхимальные стромальные клетки в процессах роста и reparации тканей. *Acta naturae.* Т. 3. № 4. С. 32. (*Kalinina N.I., Sysoeva V.Yu., Rubina K.A., Parfenova E.V., Tkachuk.* 2011. Mesenchymal stem cells in the processes of tissue growth and repair. *Acta naturae.* V. 3. № 4. P. 32.)
- Курьянов П.С., Разуваев А.С., Вавилов В.Н.* 2008. Гиперплазия интимы в зоне сосудистого анастомоза. Ангиология и сосудистая хирургия. Т. 14. № 4. С. 146. (*Kuryanov P.S., Razuvaev A.S., Vavilov V.N.* 2008. Intimal hyperplasia in the area of vascular anastomosis. *Angiol. Vascular Surgery.* V. 14. № 4. P. 146.)
- Попов Г.И., Крюков А.Е., Попрядухин П.В., Нащекина Ю.А., Иванькова Е.М., Вавилов В.Н., Юдин В.Е., Смирнова Н.В.* 2018. Выбор оптимальных методов посева и культивирования мезенхимальных стволовых клеток на биоразлагаемой матрице из L-полилактида. *Цитология.* Т. 60. № 4. С. 279. (*Popov G.I., Kryukov A.E., Popryadukhin P.V., Naschekina Yu.A., Ivankova E.M., Vavilov V.N., Yudin V.E., Smirnova N.V.* 2018. Determining optimal methods of cell seeding and cultivation on L-polylactide biodegradable scaffold. *Tsitolgiya.* V. 60. № 4. P. 279.)
- Acland R.D.* 1972. Signs of patency in small vessel anastomosis. *Surgery.* V. 72. P. 744.
- Hashi C.K., Zhu Y., Yang G., Young W.L., Hsiao B.S., Wang K., Chu B., Li S.* 2007. Antithrombogenic property of bone marrow mesenchymal stem cells in nanofibrous vascular grafts. *Proc. Natl. Acad. Sci.* V. 104. P. 11915.
- Krawieca J.T., Vorp D.A.* 2012. Adult stem cell-based tissue engineered blood vessels: A review. *Biomaterials.* V. 33. P. 3388.
- Popryadukhin P.V., Popov G.I., Yukina G.Yu., Dobrovolskaya I.P., Ivan'kova E.M., Vavilov V.N., Yudin V.E.* 2017. Tissue-engineered vascular graft of small diameter based on electro-spun polylactide microfibers. *Int. J. Biomater.* Article ID 9034186. <https://doi.org/10.1155/2017/9034186>
- Row S., Santandreu A., Swartz D.D., Andreadis S.T.* 2017. Cell-free vascular grafts: Recent developments and clinical potential. *Technology (Singap. World Sci.).* V. 5. P. 13.
- Sullivan S.J., Brockbank K.G.M.* 2000. Small-diameter vascular grafts. In: *Principles of tissue engineering.* Second edition. Acad. Press. P. 447.
- Villalona G.A., Udelesman B., Duncan D.R., McGillicuddy E., Sawh-Martinez R.F., Hibino N., Painter C., Mirensky T., Erickson B., Shinoka T., Breuer C.K.* 2010. Cell-seeding techniques in vascular tissue engineering. *Tiss. Eng. Part B. Rev.* V. 16. P. 341.
- Zhu Y., Carido M., Meinhardt A., Kurth T., Karl M.O., Ader M., Tanaka E.M.* 2013. Three-dimensional neuroepithelial culture from human embryonic stem cells and its use for quantitative conversion to retinal pigment epithelium. *PLoS One.* V. 8. P. e54552. <https://doi.org/10.1371/journal.pone.0054552>

Evaluation of Tissue-Engineered Vascular Grafts Based on the Biodegradable Scaffold and Mesenchymal Stem Cells in the Long-Term *in vivo* Experiment

G. I. Popov^{a,*}, V. N. Vavilov^a, P. V. Popryadukhin^b, G. Yu. Yukina^a, E. G. Sukhorukova^a, A. A. Davydov^a, E. M. Ivankova^b, A. E. Kryukov^b, N. V. Smirnov^{b,c}, and V. E. Yudin^{b,c}

^aPavlov First Saint Petersburg State Medical University, St. Petersburg, 197022 Russia

^bInstitute of Macromolecular Compounds, Russian Academy of Sciences, St. Petersburg, 199004 Russia

^cPeter the Great St. Petersburg State Polytechnic University, St. Petersburg, 195251 Russia

*e-mail: trek-4300@yandex.ru

In order to create a tissue-engineered vascular graft, the bioresorbable tubular scaffold from poly(L-lactide) and the method for mesenchymal stem cells seeding and culturing were developed. In group 1, scaffolds without pre-seeded cells were implanted into the abdominal rat aorta; in group 2 scaffolds with pre-cultured rat adipose-derived mesenchymal cells were used. Follow-up in the group 1 was from 2 days to 64 weeks ($n = 36$), in the group 2 up to 72 weeks ($n = 42$). Patency in the group 1 – 86% (31/36), in the group 2 – 97% (41/42). Histology revealed gradual polymer biodegradation and replacement with connective tissue, however, after 64 weeks a total polymer resorption was achieved and an aneurysm formation was noted in all cases in the group 1. Cells amount and neoadventitia thickness were above in the group 2, however the graft morphology did not fundamentally differ from the group 1. Aneurysm formation was noted only in 45% of grafts at the later stages of follow-up in the group 2. Thus, the preliminary cultivation of mesenchymal stem cells on the scaffold led to a pronounced host cellular reaction, which probably made it possible to reduce the incidence of aneurysm formation, but did not ensure the formation of a natural vessel structure.

Keywords: tissue engineering, artificial vessel, bioresorbable scaffold, poly(L-lactide), adipose-derived mesenchymal stem cells